



# On Semi-Essential Subsemimodules in Multiplication Semimodules

Pritam Gujarathi <sup>1\*</sup> and Kishor Pawar <sup>2</sup>

<sup>1</sup> Department of Engineering Sciences,  
Ramrao Adik Institute of Technology (RAIT), Nerul, Navi Mumbai – 400 706

pritam.gujarathi@gmail.com

<sup>2</sup> Department of Mathematics, School of Mathematical Sciences,  
North Maharashtra University, Jalgaon – 425 001

kfpawar@nmu.ac.in

## ABSTRACT

The Semi essential subsemimodule was defined in [4]. In this paper we generalize some results of semi essential submodule to semi essential subsemimodules in multiplication semimodule.

## Indexing terms/Keywords

Semiring; semimodule; essential subsemimodule; semi-essential subsemimodule.

## SUBJECT CLASSIFICATION

2010 AMS Classification 2010: 16Y60.

## 1. INTRODUCTION

In [4]-[7] the author has investigated and studied different classes of essential ideals and essential subsemimodules. The notion of semi-essential subsemimodule was introduced in [6] by Kishor Pawar and Pritam Gujarathi. In this paper we generalize the properties of semi-essential module over multiplication semimodule [1] and also we give conditions when an  $R$ -subsemimodule of multiplication subsemimodule becomes semi-essential subsemimodule.

## 2. PRELIMINARIES

**Definition 2.1. [3]** A semiring is a set  $R$  together with two binary operations called addition  $(+)$  and multiplication  $(\cdot)$  such that  $(R, +)$  is a commutative monoid with identity element  $0_R$ ;  $(R, \cdot)$  is a monoid with identity element  $1$ ; multiplication distributes over addition from either side and  $0$  is multiplicative absorbing, that is,  $a \cdot 0 = 0 \cdot a = 0$  for each  $a \in R$ . A semiring  $R$  is said to have a unity if there exists  $1_R \in R$  such that  $1_R \cdot a = a \cdot 1_R = a$  for each  $a \in R$ .

**For e.g.:** The set  $\mathbb{N}$  of non-negative integers with the usual operations of addition and multiplication of integers is a semiring with  $1_{\mathbb{N}}$ .

**Definition 2.2. [3]** Let  $R$  be a semiring. A left  $R$ -semimodule is a commutative monoid  $(M, +)$  with additive identity  $0_M$  for which we have a function  $R \times M \rightarrow M$  defined by  $(r, m) \mapsto r \cdot m$  and called scalar multiplication which satisfies the following conditions for all  $r$  and  $r'$  of  $R$  and all elements  $m$  and  $m'$  of  $M$ ,

1.  $(r \cdot r')m = r(r' \cdot m)$
2.  $r \cdot (m + m') = r \cdot m + r \cdot m'$
3.  $(r + r') \cdot m = r \cdot m + r' \cdot m$
4.  $1_R \cdot m = m$  (If exists)
5.  $r \cdot 0_M = 0_M = 0_R \cdot m$ .

**Convention:** In this paper all semirings considered will be assumed to be commutative semirings with unity.

**Definition 2.3 [2]:** Let  $R$  be a semiring and  $M$  be an  $R$ -semimodule. A subsemimodule  $N$  of  $M$  is called prime if

- i)  $N$  is proper subsemimodule of  $M$  and



- ii) If for any  $m \in M, r \in R, mr \in N \Rightarrow m \in N$  or  $r \in A_N(M) = \{a \in R \mid aM \subseteq N\}$ .

**Definition 2.4 [3]:** A nonzero  $R$ -subsemimodule  $N$  of  $M$  is called semi-essential if  $N \cap P \neq 0$  for each nonzero prime  $R$ -subsemimodule  $P$  of  $M$ .

### 3. SEMI-ESSENTIAL SUBSEMIMODULES IN MULTIPLICATION SEMIMODULES

In this section, we give a condition under which an  $R$ -subsemimodule  $N$  of a faithful multiplication  $R$ -semimodule  $M$  becomes semi essential.

**Definition 3.1.[3]** An  $R$ -semimodule  $M$  is called a multiplication semimodule where  $N$  is a subsemimodule of  $M$ , then there exists an ideal  $I$  of  $R$  such that

$$N = IM. I = (N : M) = \{r \in R / rm \subseteq N\}$$

**Proposition 3.2[6]:** A nonzero  $R$ -subsemimodule  $N$  of  $M$  is semi-essential if and only if for each nonzero prime  $R$ -subsemimodule  $P$  of  $M$  there exists  $x \in P$  and there exists  $r \in R$  such that  $0 \neq rx \in N$ .

**Proposition 3.3[6]:** Let  $M$  be an  $R$ -subsemimodules and let  $N_1$  and  $N_2$  be  $R$ -subsemimodules of  $M$  such that  $N_1$  is an  $R$ -subsemimodules of  $N_2$ . If  $N_1$  is a semi-essential  $R$ -subsemimodule of  $M$ , then  $N_2$  is a semi-essential  $R$ -subsemimodule of  $M$ .

**Corollary 3.4 [6]:** Let  $N_1$  and  $N_2$  are  $R$ -subsemimodules of  $M$ . If  $N_1 \cap N_2$  is a semi-essential  $R$ -subsemimodule of  $M$ , then  $N_1$  and  $N_2$  are semi-essential.

**Proposition 3.5[6]:** Let  $N_1$  and  $N_2$  are  $R$ -subsemimodules of  $M$  such that  $N_1$  is essential and  $N_2$  is semi-essential. Then  $N_1 \cap N_2$  is a semi-essential  $R$ -subsemimodule of  $M$ .

**Lemma 3.6 [6]:** Let  $N$  be an  $R$ -subsemimodule of  $M$  and let  $P$  be a prime subsemimodule of  $M$ . If  $(N \cap P : x) = \text{ann}(M)$ , for each  $x \in M$  and  $x \notin N \cap P$ , then  $N \cap P$  is a prime  $R$ -subsemimodule of  $M$ .

**Theorem 3.7 [8]** If  $M$  is finitely generated multiplication semimodule over a semiring  $R$ ,  $P$  is a strong  $k$ -ideal of  $R$  containing  $\text{ann}(M)$ , then  $PM$  is a prime subsemimodule of  $M$ .

**Theorem 3.8.** Let  $M$  be a faithful multiplication  $R$ -semimodule and  $N$  is an  $R$ -subsemimodule of  $M$  such that  $N = IM$  for some ideal  $I$  of  $R$ . If  $N$  is semi essential if and only if  $I$  is semi essential with  $I \cap P = 0$ , where  $P$  is strong prime  $k$ -ideal of  $R$  containing  $\text{ann}(M)$ .

**Proof** Since  $M$  is faithful multiplication  $R$ -semimodule then  $(I \cap P)M = 0$  Implies  $IM \cap PM = 0$ .

$PM$  is prime  $R$ -subsemimodule of  $M$  and  $N = IM$  is semiessential  $R$  subsemimodule of  $M$  therefore  $PM = 0$ . Implies  $P = 0$ . Hence  $I$  is semiessential ideal of  $R$ .

Conversely, Let  $N \cap P = 0$ , where  $P$  is non zero prime  $R$ -subsemimodule of  $M$ . Since  $M$  is multiplication semimodule there exists a strong prime  $k$ -ideal  $P'$  of  $R$  such that  $P = P'M$ . Hence  $N \cap P = IM \cap P'M = (I \cap P')M = 0$ . But  $M$  is faithful implies  $I \cap P' = 0$ . Since  $I$  is semi essential ideal of  $R$ , then  $P' = 0$ . Therefore  $P = 0$  implies  $N$  is semiessential  $R$  subsemimodule of  $M$ .

**Theorem 3.9:** Let  $M$  be a faithful multiplication  $R$ -semimodule Then  $N$  is a semiessential  $R$ -subsemimodule of  $M$  if and only if  $(N : x)$  is a semi-essential ideal of  $R$  for each  $x \in M$

**Proof:** Suppose that  $N$  is semi essential. By above Theorem 2.7  $M$  is faithful multiplication of  $R$  semimodule then  $(N : M)$  is semi essential  $k$ -ideal of  $R$ . But  $(N : M) \subseteq (N : x)$ . Therefore for each  $x \in M, N = (N : M)M \subseteq (N : x)M$ . Implies  $(N : x)M$  is semi essential  $R$ -subsemimodule of  $M$

And consequently  $(N : x)$  is a semiessential  $k$ -ideal of  $R$ .



Conversely assume that  $(N : x)$  is semi essential  $k$ -ideal of  $R$  for each  $x \in M$ . Let  $P$  be a nonzero prime  $R$ -submodule of  $M$  and let  $0 \neq y \in P$ . Thus  $(N : y)$  is semi essential. Since  $M$  is multiplication then  $P = P'M$ , where  $P$  is a strong  $k$ -prime ideal of  $R$ . Hence  $(N : y) \cap P' \neq 0$ . By assumption  $M$  is faithful, So  $(N : x)M \cap P'M \neq 0$ . Thus  $N \cap P \neq 0$ .

**Proposition 3.10:** Let  $M$  be a faithful multiplication  $R$ -semimodule and let  $N$  be nonzero prime  $R$ -subsemimodule of  $M$ . If  $N$  is not minimal prime, then  $N$  is semiessential.

**Proof:** Since  $M$  is multiplication and  $N$  is prime, then there exists a strong prime  $k$ -ideal  $P'$  of  $R$  such that  $\text{ann}(M) \subseteq P'$  such that  $N = P'M$ . Let  $P$  be nonzero prime  $R$ -subsemimodule of  $M$  such that  $N \cap P = 0$ . Since  $N$  is not minimal prime there exists a minimal prime  $R$ -subsemimodule  $L$  of  $M$  such that  $L \subset N$ . Thus there exists a strong minimal prime ideal  $P''$  of  $R$  such that  $\text{ann}(M) \subseteq P''$  and  $L = P''M \neq M$ . Rest of the proof is same as in ring.

## REFERENCES

1. Ali S. Mijbass and Nada K. Abdullah, Semi-Essential Submodules and Semi-Uniform Modules, Journal of Kirkuk University-Scientific Studies, Vol. 4, No.1, 48-58, (2009).
2. Gursel Yesilot and Kursat Hakan Oral, On Prime Subsemimodules of Semimodules, International Journal of Algebra, Vol. 4, no. 1, 53 – 60, (2010).
3. J.S. Golan, Semirings and their Applications, Kluwer Academic Publishers, (1992).
4. K. F. Pawar and R. P. Deore, On Essential Ideal and Radical Class, Int. J. of Pure and Appl. Math. Sci. Vol. 5(1), 1-5, (2012).
5. Kishor Pawar, A Note on Essential Subsemimodule, New Trends in Mathematical Sciences, Vol. 1(2), 18-21, (2013).
6. Kishor Pawar and Pritam Gujarathi, On Semi-Essential Subsemimodules, Journal of Advances in Mathematics Vol. 12, No. 1, pp. 5805-5807, (2015).
7. Kishor Pawar, On Weak Essential Ideals of Semiring, Communications in Mathematics and Applications, Vol. 6(1), 17-20, (2015).
8. S. Ebrahimi Atani and M. Shajari Kohan, A Note on  $\pi$ -nitely generated multiplication semimodules over commutative semirings, International Journal of Algebra, Vol. 4, no. 8, 389-396, (2010).
9. S. Ebrahimi Atani and S. Khojasteh G. Ghaleh, On Multiplication Semimodules, International Mathematical Forum, 1, no. 24, 1175-1180, (2006).
10. Dutta, T.K. and Das, M.L. On Strongly Prime Semiring, Bull. Malays Math. Sci. Soc, 30(2), 135-141, (2007).